

MONOCLONAL ANTIBODY

Biotin labeled Anti-mouse Sca-1

Code No.	Clone	Subclass	Quantity	Concentration
D052-6	238B	Rat IgG2a	100 µL	1 mg/mL

BACKGROUND: Sca-1 is a member of the Ly-6 antigen family which molecular mass of 8 kDa under nonreducing conditions and of 18 kDa under reducing conditions. Mouse hematopoietic stem cell expresses low levels of Thy-1 antigen (Thy-1^{lo}) and to be lineage-negative (Lin⁻); not express markers characteristic of B cells (B220), granulocytes (Gr-1), myelomonocytic cells (Mac-1) and T lymphocytes (CD4 and CD8). Recently, new monoclonal antibody, anti-Sca-1, was used to purify stem cells from the Thy-1^{lo}, Lin⁻ bone marrow subpopulation. Thy-1^{lo}, Lin⁻, Sca-1⁺ (but not the Thy-1^{lo}, Lin⁻, Sca-1⁻) population of bone marrow cells are highly purified pluripotent stem cells. They read out with nearly unit efficiency in assays for primitive myeloerythroid and thymic progenitor, and have a capability to admit lethally irradiated mouse to survive and be restored in all blood-cell lineages. The Thy-1^{lo}, Lin⁻, Sca-1⁺ subpopulation thought to have all stem cells present in the bone marrow.

SOURCE: This antibody was purified from culture supernatant using protein G agarose. This hybridoma (clone 238B) was established by fusion of mouse myeloma cell P3X with rat splenocyte immunized with mouse Sca-1 transfected LO cells.

FORMULATION: 100 µg IgG in 100 µL volume of PBS containing 1% BSA and 0.09% NaN₃.

*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at 4°C.

REACTIVITY: This antibody reacts with mouse Sca-1 on Flow cytometry.

APPLICATION:

Flow cytometry: 10 µg/mL (final concentration)

*Please refer to the data sheet (MBL code no. D052-3) for other applications.

Detailed procedure is provided in the following **PROTOCOL.**

INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.

RELATED PRODUCTS:


- D052-3 Anti-mouse Sca-1 (238B)
- D052-5 PE labeled anti-mouse Sca-1 (238B)
- D160-3 Anti-mouse Ly49Q (2E6)
- D160-4 FITC labeled anti-mouse Ly49Q (2E6)
- D160-5 PE labeled anti-mouse Ly49Q (2E6)

SPECIES CROSS REACTIVITY:

Species	Human	Mouse	Rat
Cell	Not Tested	LO	Not Tested
Reactivity on FCM		+	

REFERENCE:

- 1) Petersen, B. E., *et al.*, *Hepatology* **37**, 632-640 (2003)



Flow cytometric analysis of Biotin labeled mouse Sca-1 expression on LO cells. Open histogram indicates the reaction of Isotypic control to the cells. Shaded histogram indicates the reaction of D052-6 to the cells.

PROTOCOL:

Flow cytometric analysis for adherent cells

We usually use Fisher tubes or equivalents as reaction tubes for all step described below.

- 1) Detach the cells from culture dish by cell dissociation buffer.
- 2) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.1% NaN₃].
- 3) Resuspend the cells with washing buffer (5x10⁶ cells/mL).
- 4) Add 50 µL of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 5) Add 10 µL of normal goat serum containing 1 mg/mL

normal human IgG and 0.1% NaN₃ to the cell pellet after tapping. Mix well and incubate for 10 minutes at room temperature.

- 6) Add 40 µL of the Biotin labeled anti-mouse Sca-1 monoclonal antibody (238B) (25 µg/mL) diluted with the washing buffer. Mix well, and incubate for 30 minutes at room temperature.
- 7) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 8) Add 20 µL of 1:40 FITC conjugated streptavidin (MBL; code no. IM-0307) diluted with the washing buffer. Mix well and incubate for 15 minutes at room temperature.
- 9) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 10) Resuspend the cells with 500 µL of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; LO)