

MONOCLONAL ANTIBODY

Anti-Mouse TLR4-MD-2 complex

Code No.	Clone	Subclass	Quantity	Concentration
D079-3	MTS510	Rat IgG2a κ	100 μ L	1 mg/mL

BACKGROUND: Toll, a Drosophila receptor molecule with extracellular leucine-rich repeat (LRR), has a role in triggering innate defenses against bacteria or fungi. TLR4 (Toll-like receptor 4) is a member of TLR family, which is human homologue of Toll protein. It has extracellular LRR and an intracellular signaling domain, which is similar to the type I IL-1 receptor. TLR4 is expressed in subpopulations of cells including myeloid cells, B cells, monocytes, and endothelial cells. Recent studies have suggested that TLR4 might act as a receptor for LPS (lipopolysaccharide). TLR4 alone is not capable of sensing and signaling the presence of LPS, but another molecule MD-2, which is physically associated with TLR4, is required for LPS recognition through TLR4.

SOURCE: This antibody was purified from culture supernatant using protein G agarose. This hybridoma was established by fusion of mouse myeloma cell SP2/0 with Wister rat lymph node cells immunized with mouse Ba/F3 cells expressing mouse TLR4-MD-2 complex.

FORMULATION: 100 μ g IgG in 100 μ L volume of PBS containing 50% glycerol. No preservative is contained.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at -20°C.

REACTIVITY: This antibody reacts with mouse TLR4-MD-2 complex on Flow cytometry. It doesn't react with mouse TLR4 alone. Clone MTS510 inhibits the TNF- α production from peritoneal macrophages by LPS stimulation*.

*It is reported in reference number 12).

APPLICATIONS:

Western blotting; Not tested

Immunoprecipitation; Not tested*

Neutralization; Not tested*

*It is reported that clone MTS510 can be used in these applications in the reference number 12).

Immunohistochemistry; Not tested


Immunocytochemistry; Not tested

Flow cytometry; 10-20 μ g/mL (final concentration)

Detailed procedure is provided in the following **PROTOCOL**.

SPECIES CROSS REACTIVITY:

Species	Human	Mouse		Rat
Cells	Not Tested	Ba/F3 cells expressing mouse TLR4-MD-2	Ba/F3 cells expressing mouse TLR4 alone	Not Tested
Reactivity on FCM		+	-	



Flow cytometric analysis of TLR4-MD-2 complex expression on Ba/F3 parental cells (right) or mouse TLR4 and MD-2 co-transfected Ba/F3 cells (left). Open histogram indicates the reaction of isotypic control to the cells. Shaded histograms indicate the reaction of D079-3 to the cells.

INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.

REFERENCES:

- 1) Wakabayashi, Y., *et al.*, *J. Immunol.* **177**, 1772-1779 (2006)
- 2) Kobayashi, M., *et al.*, *J. Immunol.* **176**, 6211-6218 (2006)
- 3) Akashi-Takamura, S., *et al.*, *J. Immunol.* **176**, 4244-4251 (2006)
- 4) Saitoh, S., *et al.*, *Int. Immunol.* **16**, 961-969 (2004)
- 5) Akashi, S., *et al.*, *J. Exp. Med.* **198**, 1035-1042 (2003)
- 6) Yazawa, N., *et al.*, *Blood* **102**, 1374-1380 (2003)
- 7) Kawasaki, K., *et al.*, *J. Immunol.* **170**, 413-420 (2003)
- 8) Nagai, Y., *et al.*, *Nat. Immunol.* **3**, 667-672 (2002)
- 9) Akashi, S., *et al.*, *Int. Immunol.* **13**, 1595-1599 (2001)
- 10) Ogata, H., *et al.*, *J. Exp. Med.* **192**, 23-30 (2000)
- 11) Nomura, F., *et al.*, *J. Immunol.* **164**, 3476-3479 (2000)
- 12) Akashi, S., *et al.*, *J. Immunol.* **164**, 3471-3475 (2000)
- 13) Akashi, S., *et al.*, *Biochem. Biophys. Res. Commun.* **268**, 172-177 (2000)
- 14) Shimazu, R., *et al.*, *J. Exp. Med.* **189**, 1777-1782 (1999)

Clone MST510 is used in reference number 1) - 12).

PROTOCOL:

Flow cytometric analysis for floating cells

We usually use Fisher tubes or equivalents as reaction tubes for all steps described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.1% NaN₃].
- 2) Resuspend the cells with washing buffer (5x10⁶ cells/mL).
- 3) Add 50 µL of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 4) Add 20 µL of normal goat serum containing 1 mg/mL normal human IgG and 0.1% NaN₃ to the cell pellet after tapping. Mix well and incubate for 5 minutes at room temperature.
- 5) Add 40 µL of the primary antibody at the concentration as suggest in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Add 30 µL of 1:40 FITC conjugated anti-rat IgG (MBL; code no. IM-0827) diluted with the washing buffer. Mix well and incubate for 15 minutes at room temperature.
- 8) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 9) Resuspend the cells with 500 µL of the washing buffer and analyze by a flow cytometer.

RELATED PRODUCTS:

- D210-3 anti-TLR1/CD281 (GD2.F4)
- K0211-3 anti-Mouse TLR2/CD282 (mT2.7)
- K0212-3 anti-Mouse TLR2/CD282 (T2.5)
- D077-3 anti-Human TLR4/CD284 (HTA125)
- D077-4 FITC labeled anti-Human TLR4/CD284 (HTA125)
- D077-5 PE labeled anti-Human TLR4/CD284 (HTA125)
- D079-4 FITC labeled anti-Mouse TLR4-MD-2 complex (MTS510)
- D079-5 PE labeled anti-Mouse TLR4-MD-2 complex (MTS510)
- D205-3 anti-Mouse TLR4/CD284 (UT49)
- D205-4 FITC labeled anti-Mouse TLR4/CD284 (UT49)
- D206-3 anti-Mouse TLR4-MD-2 complex (UT15)
- D206-5 PE labeled anti-Mouse TLR4-MD-2 complex (UT15)
- K0213-3 anti-Mouse TLR9/CD289 (5G5)
- M081-3 Rat IgG2a Isotype control (2H3)
- M081-4 FITC labeled Rat IgG2a Isotype control (2H3)
- M081-5 PE labeled Rat IgG2a Isotype control (2H3)
- M081-8 Agarose conjugated Rat IgG2a Isotype control (2H3)