

Anti-Fc μ R (TOSO/FAIM3) (Mouse) mAb

CODE No.	D303-3
CLONALITY	Monoclonal
CLONE	#4B5
ISOTYPE	Rat IgG1 κ
QUANTITY	100 μ L, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	Mouse Fc μ R extracellular domain-Fc fusion protein
FORMULATION	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATIONS-CONFIRMED

<u>Immunocytochemistry</u>	10 μ g/mL
<u>Flow cytometry</u>	10 μ g/mL

SPECIES CROSS REACTIVITY on FCM

Species	Human	Mouse	Rat	Hamster
Cells	Transfectant	Transfectant Splenocytes* Bone marrow cells*	Not tested	Not tested
Reactivity	-	+		

*Reactivity of this clone to mouse splenocytes and bone marrow cells on FCM is described in the reference number 1).

Entrez Gene ID 69169 (Mouse)

- REFERENCES**
- 1) Shima, H., *et al.*, *Int. Immunol.* **22**, 149-156 (2010) [FCM]
 - 2) Kubagawa, H., *et al.*, *J. Exp. Med.* **206**, 2779-2793 (2009)
 - 3) Pallasch, C. P., *et al.*, *Blood* **112**, 4213-4219 (2008)
 - 4) Proto-Siqueira, R., *et al.*, *Blood* **112**, 394-397 (2008)
 - 5) Hitoshi, Y., *et al.*, *Immunity* **8**, 461-471 (1998)




RELATED PRODUCTS

For more information, please visit our web site <https://ruo.mbl.co.jp/>.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

Immunocytochemistry

- 1) Spread the cells in the nutrient condition on a glass slide, then incubate in a CO₂ incubator overnight.
- 2) Remove the culture supernatant by careful aspiration.
- 3) Fix the cells by immersing the slide in 4% paraformaldehyde (PFA)/PBS for 10 min. at room temperature (20~25°C).
- 4) Prepare a wash container such as a 500 mL beaker with a magnetic stirrer. Then wash the fixed cells on the glass slide by soaking the slide with a plenty of PBS in the wash container for 5 min. Take care not to touch the cells. Repeat another wash once more.
- 5) Immerse the slide in 0.2% Triton X-100/PBS for 20 min. at room temperature.
- 6) Wash the slide in a plenty of PBS as in the step 4).
- 7) Add 200 µL of the primary antibody diluted with 2% fetal calf serum (FCS)/PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hr. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 8) Wash the slide in a plenty of PBS as in the step 4).
- 9) Add 200 µL of 1:500 Alexa Fluor[®] 594 conjugated anti-rat IgG (Thermo Fisher Scientific, code no. A-11007) diluted with PBS onto the cells. Incubate for 30 min. at room temperature. Keep out light by aluminum foil.
- 10) Wash the slide in a plenty of PBS as in the step 4).
- 11) Add 200 µL of 0.5 µg/mL Anti-HA-tag mAb-Alexa Fluor[®] 488 (MBL, code no. M180-A48) diluted with PBS onto the cells. Incubate for 30 min. at room temperature. Keep out light by aluminum foil.
- 12) Wash the slide in a plenty of PBS as in the step 4).
- 13) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 14) Counter stain with DAPI for 5 min. at room temperature.
- 15) Wash the slide in a plenty of PBS as in the step 4).
- 16) Promptly add mounting medium onto the slide, then put a cover slip on it.




Immunocytochemical detection of HA-tagged mouse Fc μ R/TOSO/FAIM3 in HeLa

Red: D303-3
Green: Anti-HA-tag mAb-Alexa Fluor[®] 488
(MBL, code no.M180-A48)
Blue: DAPI

Flow cytometric analysis for adherent cells

- 1) Detach the cells from culture dish.
- 2) Wash the cells once with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 3) Resuspend the cells with washing buffer (3×10^6 cells/mL).
- 4) Add 100 μ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 min. at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 5) Add 10 μ L of Clear Back (human Fc receptor blocking reagent, MBL, code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 10 min. at room temperature.
- 6) Add 30 μ L of the primary antibody at the concentration as suggested in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 20 min. at 4°C.
- 7) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration. Repeat another wash once more.
- 8) Add 30 μ L of 1:100 FITC conjugated anti-rat IgG diluted with the washing buffer. Mix well and incubate for 20 min. at room temperature.
- 9) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration.
- 10) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.



Flow cytometric detection of HA-tagged mouse Fc μ R/TOSO/FAIM3 in HeLa

Open: D303-3

Closed: Isotype control (MBL, code no. M080-3)