

Anti-Dab1 (Mouse) mAb

CODE No.	D355-3
CLONALITY	Monoclonal
CLONE	4E12
ISOTYPE	Rat IgG2a κ
QUANTITY	100 μ L, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	Mouse Dab1 (recombinant)
FORMURATION	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATION-CONFIRMED

Western blotting 1 μ g/mL for chemiluminescence detection system

APPLICATIONS-REPORTED

Immunocytochemistry Reference 1)

Immunohistochemistry Reference 1)

SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Hamster
Sample	Not tested	Transfectant	Not tested	Not tested
Reactivity		+		

Entrez Gene ID 13131 (Mouse)

REFERENCES

- 1) Onoue, A., *et al.*, *Neurosci. Res.* **88**, 23-27 (2014) [WB, IC, IHC-fr]
- 2) Uchida, T., *et al.*, *J. Neurosci.* **29**, 10653-10662 (2009)
- 3) Morimura, T., *et al.*, *J. Biol. Chem.* **280**, 16901-16908 (2005)

For more information, please visit our web site <http://ruo.mbl.co.jp/>


RELATED PRODUCTS

D355-3	Anti-Dab1 (Mouse) mAb (4E12)
D354-3	Anti-Dab1 (Mouse) mAb (4H11)
D351-3	Anti-Reelin (Mouse) mAb (2F3)
D223-3	Anti-Reelin (CR-50) mAb (RE-3B9(R3B9))
M067-3	Anti-Apolipoprotein E4 (Human) mAb (1F9)
M068-3	Anti-Apolipoprotein E (Human) mAb (3D12)
D273-3	Anti-ApoER2 (LA8) (Mouse) mAb (25G5)
7635	ApoE4/Pan-ApoE ELISA Kit
D352-3	Anti-Kiaa0319 (Mouse) mAb (56F3)
D353-3	Anti-Kiaa0319 (Mouse) mAb (62A3)
M081-3	Rat IgG2a (isotype control) (2H3)

SDS-PAGE & Western blotting

- 1) Boil the sample for 2 min. and centrifuge. Load 20 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (7.5% acrylamide) for electrophoresis.
- 2) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 3) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 4) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATION** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T (5 min. x 3 times)
- 7) Incubate the membrane with HRP-conjugated anti-rabbit IgG antibody diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 8) Wash the membrane with PBS-T (5 min. x 3 times).
- 9) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 3 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; Transfectant)



Western blot analysis of mouse *Dab1*

Lane 1: Mouse *Dab1*/COS-7
Lane 2: COS-7

Immunoblotted with Anti-*Dab1* (Mouse) mAb (D355-3)

The sample was kindly provided by Dr. Mitsuharu Hattori.
(Department of Biomedical Science, Graduate School of
Pharmaceutical Sciences, Nagoya City University)