

MONOCLONAL ANTIBODY

# Anti-Integrin $\alpha 7$ (Mouse) mAb-PE

<b>Code No.</b>	<b>Clone</b>	<b>Subclass</b>	<b>Quantity</b>
K0046-5	3C12	Mouse IgG1	1 mL (50 tests)

**BACKGROUND:** The integrin family of adhesion molecules participate in important cell-cell and cell-extracellular matrix interactions in a diverse range of biological processes. Integrins are heterodimers consisting of a  $\alpha$  subunit and  $\beta$  subunit. Both  $\alpha$  and  $\beta$  subunit are transmembrane proteins with large extracellular domains (>100 kDa for  $\alpha$  subunit and >75 kDa for  $\beta$  subunit) that interact with extracellular matrix proteins and relatively small cytoplasmic domains (50 amino acids or less, except for the  $\beta 4$  subunit) that interact with cytoskeletal proteins. The adhesiveness of integrins is dynamically regulated in response to cytoplasmic signals, termed "inside-out" signaling. It has been reported that, upon ligand binding, integrins regulate many intracellular signaling pathways that involve cytoplasmic alkalization, intracellular  $\text{Ca}^{2+}$  fluctuation, inositol lipid metabolism, protein kinase C, MAP kinase and phosphatidylinositol kinase. Integrin  $\alpha 7$  is a specific cellular receptor for the basement membrane protein laminin-1, as well as for the laminin isoforms-2 and -4. The  $\alpha 7$  subunit is expressed mainly in skeletal and cardiac muscle and may be involved in differentiation and migration processes during myogenesis. Absence of integrin  $\alpha 7$  results in muscular dystrophy is revealed.

**SOURCE:** This antibody was purified from hybridoma (3C12) supernatant using protein A agarose. This hybridoma was established by fusion of mouse myeloma cell SP2/0 with Integrin  $\alpha 7$  knockout C57/B6 mouse splenocyte immunized with mouse myoblasts.

**FORMULATION:** 50 tests in 1 mL volume of PBS containing 1% BSA and 0.09%  $\text{NaN}_3$ .

\*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at 4°C.

**REACTIVITY:** This antibody reacts with mouse Integrin  $\alpha 7$  on flow cytometry.

**APPLICATIONS:**

- Western blotting; Not tested
- Immunoprecipitation; Not tested
- Immunohistochemistry; Not tested
- Immunocytochemistry; Not tested
- Flow cytometry; 20  $\mu\text{L}$  (ready for use)

Detailed procedure is provided in the following **PROTOCOL**.

**INTENDED USE:**

For research use only. Not for clinical diagnosis.

**SPECIES CROSS REACTIVITY:**


Species	Human	Mouse	Rat
Cell	Not tested	C2C12	Not tested
Reactivity on FCM		+	

**REFERENCES:**

- 1) Xynos, A., *et al. J. Cell. Sci.* **126**, 2236-2245 (2013) [FCM]
- 2) Majka, S. M., *et al. Adipocyte* **1**, 215-229 (2012) [FCM]
- 3) Mielenz, D., *et al. J. Biol. Chem.* **276**, 13417-13426 (2001)

**RELATED PRODUCTS:**

- K0046-3 Anti-Integrin  $\alpha 7$  (Mouse) mAb (3C12)
- K0046-4 Anti-Integrin  $\alpha 7$  (Mouse) mAb-FITC (3C12)
- K0047-3 Anti-Integrin  $\alpha 7$  (Mouse) mAb (6A11)
- M075-5 Mouse IgG1 (isotype control)-PE (2E12)



**Flow cytometric analysis of mouse Integrin  $\alpha 7$  expression on NIH/3T3 cells (left) and C2C12 cells (right). Open histogram indicates the reaction of isotypic control to the cells. Shaded histograms indicate the reaction of K0046-5 to the cells.**

**PROTOCOL:**

**Flow cytometric analysis for floating cells**

We usually use Fisher tubes or equivalents as reaction tubes for all step described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.09%  $\text{NaN}_3$ ].
- 2) Resuspend the cells with washing buffer ( $5 \times 10^6$  cells/mL).
- 3) Add 50  $\mu\text{L}$  of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature

(20~25°C). Remove supernatant by careful aspiration.

- 4) Add 10  $\mu$ L of normal goat serum containing 1 mg/mL normal human IgG and 0.1% NaN<sub>3</sub> to the cell pellet after tapping. Mix well and incubate for 10 minutes at room temperature.
- 5) Add the primary antibody at the amount as suggested in the **APPLICATIONS**. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.

(Positive control for flow cytometry: C2C12)