

MONOCLONAL ANTIBODY

Anti-HLA-A2 (Human) mAb-PE

Code No.	Clone	Subclass	Quantity
K0186-5	BB7.2	Mouse IgG2b	1 mL (50 tests)

BACKGROUND: Human leukocyte antigen-A2 (HLA-A2) is a class I molecule of the human major histocompatibility complex (MHC). It is a heterodimer composed of a 43 kDa α subunit that is non-covalently associated with the 12 kDa β 2-microglobulin protein. HLA-A2, like other class I molecules, binds peptides from proteins degraded in the cytoplasm and plays a role in antigen presentation and interaction with CD8⁺ T cells. HLA-A2 is the most common HLA-A allele in the Caucasian and American Indian populations (50% and ~30%, respectively) and A2 restricted T cell epitopes have been studied widely.

SOURCE: This antibody was purified from hybridoma (clone BB7.2) supernatant using protein A agarose. This hybridoma was established by fusion of mouse myeloma cell NS-1 with Balb/c mouse splenocyte immunized with papain solubilized HLA-A2.

FORMULATION: 50 tests in 1 mL volume of PBS containing 1% BSA and 0.1% ProClin 150.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at 4°C.

REACTIVITY: This antibody reacts with human HLA-A2 on Flow cytometry.

APPLICATION:

Flow cytometry; 20 μ L (Ready for use)

*Please refer to the data sheet (MBL; code no. K0186-3) for other applications.

Detailed procedure is provided in the following **PROTOCOLS**.

SPECIES CROSS REACTIVITY:

Species	Human	Mouse	Rat
Cell	T2	Not tested	Not tested
Reactivity on FCM	+		

INTENDED USE:


For Research Use Only. Not for use in diagnostic procedures.

REFERENCES:

- 1) Shen, H., *et al.*, *Cancer Immunol. Immunother.* **62**, 393-403 (2013)[FCM]
- 2) Kozako, T., *et al.*, *J. Immunol.* **177**, 5718-5726 (2006)
- 3) Yamano, Y., *et al.*, *J. Exp. Med.* **199**, 1367-1377 (2004)
- 4) Rodolfo, M., *et al.*, *Cancer Res.* **63**, 6948-6955 (2003)
- 5) Smith, M. E. F., *et al.*, *PNAS* **86**, 5557-5561 (1989)
- 6) Parham, P., *et al.*, *Human Immunology* **3**, 277-299 (1981)

Clone BB7.2 is used in these references.

As clone BB7.2 is really famous all over the world, a lot of researches have been reported. These references are a part of such reports.



Flow cytometric analysis of HLA-A2 expression on T2 cells (right) and Jurkat Cells (left). Open histogram indicate the reaction of isotypic control to the cells. Shaded histograms indicate the reaction of K0186-5 to the cells.

PROTOCOLS:

Flow cytometric analysis for floating cells

We usually use Fisher tubes or equivalents as reaction tubes for all step described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.09% NaN₃].
*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.
- 2) Resuspend the cells with washing buffer (5x10⁶ cells/mL).
- 3) Add 50 μ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 4) Add 20 μ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 minutes at room temperature.

- 5) Add 40 μ L of the primary antibody at the concentration of as suggest in the **APPLICATIONS** diluted with the washing buffer. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.

K0125-3 Anti-HLA-G (Human) mAb (MEM-G/1)
K0019-1 Anti-HLA-DR (Human) mAb (LN-3)
M077-5 Mouse IgG2b (isotype control)-PE (3D12)

(Positive control for Flow cytometry; T2)

Flow cytometric analysis for whole blood cells

We usually use Fisher tubes or equivalents as reaction tubes for all step described below.

- 1) Add 50 μ L of the primary antibody at the concentration of as suggest in the **APPLICATIONS** diluted with the washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.09% NaN₃] into each tube.
- 2) Add 50 μ L of whole blood into each tube. Mix well, and incubate for 30 minutes at room temperature (20~25°C).
- 3) Add 1 mL of washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 4) Lyse with OptiLyse C (for analysis on Beckman Coulter instruments) or OptiLyse B (for analysis on BD instruments), using the procedure recommended in the respective package inserts.
- 5) Add 1 mL of H₂O to each tube and incubate for 10 minutes at room temperature.
- 6) Centrifuge at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Add 1 mL of washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 8) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.

RELATED PRODUCTS:

K0186-3 Anti-HLA-A2 (Human) mAb (BB7.2)
K0186-4 Anti-HLA-A2 (Human) mAb-FITC (BB7.2)
K0186-5 Anti-HLA-A2 (Human) mAb-PE (BB7.2)
K0208-3 Anti-HLA-A24 (Human) mAb (17A10)
K0208-4 Anti-HLA-A24 (Human) mAb-FITC (17A10)
K0208-5 Anti-HLA-A24 (Human) mAb-PE (17A10)
K0208-A48 Anti-HLA-A24 (Human) mAb
-Alexa Fluor[®] 488 (17A10)
K0208-A64 Anti-HLA-A24 (Human) mAb
-Alexa Fluor[®] 647 (17A10)
K0209-3 Anti-HLA-A24 (Human) mAb (22E1)
K0209-4 Anti-HLA-A24 (Human) mAb-FITC (22E1)
K0209-5 Anti-HLA-A24 (Human) mAb-PE (22E1)
D226-3 Anti-HLA class I (HLA-A,B,C) (Human) mAb
(EMR8-5)
K0126-3 Anti-HLA-E (Human) mAb (MEM-E/02)
K0215-3 Anti-HLA-E (Human) mAb (4D12)