

Smart-IP Series

Anti-Myc-tag mAb-Magnetic Agarose

CODE No.	M047-10
CLONALITY	Monoclonal
CLONE	PL14
ISOTYPE	Mouse IgG1 κ
QUANTITY	20 tests (Gel: 200 μ L)
SOURCE	Purified IgG from mouse ascites fluid
IMMUNOGEN	6myc-tagged fusion protein
FORMULATION	400 μ g of antibody is covalently coupled to 200 μ L of magnetic agarose gel and provided as 400 μ L gel slurry suspended in PBS/0.1% ProClin 150
STORAGE	This gel slurry is stable for one year from the date of purchase when stored at 4°C.
APPLICATION-CONFIRMED	
<u>Immunoprecipitation</u>	10 μ L of gel/400 μ L of cell extract from 2×10^6 cells
REFERENCE	1) Bae, S. J., <i>et al.</i> , <i>J. Cell Biochem.</i> 118 , 2219-2230 (2017) [IP]


RELATED PRODUCTS

Please visit our website at <https://ruo.mbl.co.jp/>

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

Immunoprecipitation

- 1) Wash 2×10^6 cells 3 times with PBS and suspend them in 400 μL of cold Lysis buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40], then sonicate briefly (up to 10 sec.).
- 2) Centrifuge the tube at 12,000 x g for 5 min. at 4°C and transfer the supernatant to another tube.
- 3) Add magnetic beads as suggested in the **APPLICATION** into 400 μL of the cell lysate. Mix well and incubate with gentle agitation for 30 min. at 4°C.
- 4) Place the tube on the magnetic rack (MBL; code no. 3190) for a few seconds.
- 5) Remove the supernatant.
- 6) Add 1 mL of cold Lysis buffer and resuspend the magnetic beads.
- 7) Place the tube on the magnetic rack for a few seconds.
- 8) Remove the supernatant.
- 9) Repeat Steps 6)-8) 3 times.
- 10) Resuspend the magnetic beads in 50 μL of Laemmli's sample buffer, boil for 3 min., and place the tube on the magnetic rack for a few seconds.
- 11) Load 10 μL of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) and carry out electrophoresis.
- 12) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 13) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 14) Incubate the membrane with 1:10,000 of Anti-Myc-tag mAb-HRP-Direct (MBL; code no. M192-7) diluted with 1% skimmed milk (in PBS, pH 7.2) PBS for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 15) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 16) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 17) Expose to an X-ray film in a dark room for 30 sec. Develop the film as usual settings. The condition for exposure and development may vary.



Immunoprecipitation of Myc-tagged protein

Lane 1: Parental cell (293T)

Lane 2: Myc-tagged protein in 293T

Immunoblotted with Anti-Myc-tag mAb-HRP-Direct (MBL; code no. M192-7)