

 **My select** sampler set

# Rat IgG2b (isotype control)-Alexa Fluor<sup>®</sup> 647

Code No.	Clone	Subclass	Quantity	Concentration
M090-A64MS	3G8	Rat IgG2b $\kappa$	20 $\mu$ L	1 mg/mL

**SOURCE:** This antibody was purified from hybridoma (clone 3G8) supernatant using protein G agarose. This hybridoma was established by fusion of mouse myeloma cell P3U1 with rat lymph nodes immunized with KLH.

**FORMULATION:** 20  $\mu$ g IgG in 20  $\mu$ L volume of PBS containing 1% BSA and 0.09% NaN<sub>3</sub>.

\*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at 4°C.

**REACTIVITY:** No specific binding is detected on mouse splenocytes.

**APPLICATION:**

Flow cytometry: This antibody can be used as a negative isotypic control. The concentration will depend on the conditions.

Detailed procedure is provided in the following **PROTOCOL.**

**INTENDED USE:**


For Research Use Only. Not for use in diagnostic procedures.

**PROTOCOL:**

**Flow cytometric analysis for floating cells**

We usually use Fisher tubes or equivalents as reaction tubes for all steps described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.1% NaN<sub>3</sub>].
- 2) Resuspend the cells with washing buffer (6x10<sup>6</sup> cells/mL).
- 3) Add 50  $\mu$ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 4) Add 10  $\mu$ L of normal goat serum to the cell pellet after tapping. Mix well and incubate for 10 minutes at room temperature.
- 5) Add the isotype control antibody at the concentrations comparable to those of the specific antibody of interest. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.



**Flow cytometric analysis of rat IgG2b reactivity of mouse splenocytes.**

## RELATED PRODUCTS:

### Functional grade antibodies

- M080-3M2 Rat IgG1 (isotype control) (1H5)
- M081-3M2 Rat IgG2a (isotype control) (2H3)
- M090-3M2 Rat IgG2b (isotype control) (3G8)
- M075-3M2 Mouse IgG1 (isotype control) (2E12)
- M076-3M2 Mouse IgG2a (isotype control) (6H3)
- M077-3M2 Mouse IgG2b (isotype control) (3D12)

### Purified antibodies

- M080-3 Rat IgG1 (isotype control) (1H5)
- M080-4 Rat IgG1 (isotype control)-FITC (1H5)
- M080-5 Rat IgG1 (isotype control)-PE (1H5)
- M080-A48 Rat IgG1 (isotype control)-Alexa Fluor® 488 (1H5)
- M081-3 Rat IgG2a (isotype control) (2H3)
- M081-4 Rat IgG2a (isotype control)-FITC (2H3)
- M081-5 Rat IgG2a (isotype control)-PE (2H3)
- M081-8 Rat IgG2a (isotype control)-Agarose (2H3)
- M081-9 Rat IgG2a (isotype control)-Magnetic beads (2H3)
- M081-A48 Rat IgG2a (isotype control)-Alexa Fluor® 488 (2H3)
- M090-3 Rat IgG2b (isotype control) (3G8)
- M090-4 Rat IgG2b (isotype control)-FITC (3G8)
- M090-5 Rat IgG2b (isotype control)-PE (3G8)
- M090-A48 Rat IgG2b (isotype control)-Alexa Fluor® 488 (3G8)
- M082-3 Rat IgG2c (isotype control) (6E12)
- M082-4 Rat IgG2c (isotype control)-FITC (6E12)
- M075-3 Mouse IgG1 (isotype control) (2E12)
- M075-4 Mouse IgG1 (isotype control)-FITC (2E12)
- M075-5 Mouse IgG1 (isotype control)-PE (2E12)
- M075-8 Mouse IgG1 (isotype control)-Agarose (2E12)
- M075-9 Mouse IgG1 (isotype control)-Magnetic beads (2E12)
- M075-A48 Mouse IgG1 (isotype control)-Alexa Fluor® 488 (2E12)
- M075-A64 Mouse IgG1 (isotype control)-Alexa Fluor® 647 (2E12)
- M076-3 Mouse IgG2a (isotype control) (6H3)
- M076-4 Mouse IgG2a (isotype control)-FITC (6H3)
- M076-5 Mouse IgG2a (isotype control)-PE (6H3)
- M076-9 Mouse IgG2a (isotype control)-Magnetic beads (6H3)
- M076-A48 Mouse IgG2a (isotype control)-Alexa Fluor® 488 (6H3)
- M076-A64 Mouse IgG2a (isotype control)-Alexa Fluor® 647 (6H3)
- M077-3 Mouse IgG2b (isotype control) (3D12)
- M077-4 Mouse IgG2b (isotype control)-FITC (3D12)
- M077-5 Mouse IgG2b isotype control-PE (3D12)
- M077-9 Mouse IgG2b (isotype control)-Magnetic beads (3D12)
- M077-A48 Mouse IgG2b (isotype control)-Alexa Fluor® 488 (3D12)
- M077-A64 Mouse IgG2b (isotype control)-Alexa Fluor® 647 (3D12)
- M078-3 Mouse IgG3 (isotype control) (6A3)
- M078-4 Mouse IgG3 (isotype control)-FITC (6A3)
- M079-3 Mouse IgM (isotype control) (7E10)
- M194-3 Human IgG1 isotype control chimeric mAb
- M195-3 Human IgG2 isotype control chimeric mAb
- PM035 Normal Rabbit IgG (polyclonal)
- PM035-8 Normal Rabbit IgG-Agarose (polyclonal)
- PM067 Normal Guinea Pig IgG (polyclonal)
- M189-3 Syrian Hamster IgG (isotype control)
- M199-3 Armenian Hamster IgG (isotype control)

Other related antibodies and kits are also available.  
Please visit our web site at <http://ruo.mbl.co.jp>

130603-1

## **LABEL LICENSES:**

*This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co., LTD, and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by or licensed to LIFE TECHNOLOGIES Corporation. For information on purchasing a license to this product for any other use, contact LIFE TECHNOLOGIES Corporation, Molecular Probes Detection Technologies, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.*

Alexa Fluor® is a registered trademark of Molecular Probes, Inc.