

 **My select** sampler set

Loading Control Antibody

Anti- α -Tubulin mAb

Code No.	Clone	Subclass	Quantity	Concentration
M175-3MS	2F9	Mouse IgG2a κ	20 μ L	2 mg/mL

BACKGROUND: Microtubules are one of the components of the cytoskeleton, which performs essential and diverse functions within eukaryotic cells. Microtubules are composed of a heterodimer of α and β tubulins. Tubulin is a GTP-binding protein, and extension and shortening of the microtubules are regulated by binding/hydrolysis of GTP.

SOURCE: This antibody was purified from hybridoma (clone 2F9) supernatant using protein A agarose. This hybridoma was established by fusion of mouse myeloma cell P3U1 with C3H mouse lymphocyte immunized with the synthetic peptide corresponding to N-terminal of human α -Tubulin.

FORMULATION: 40 μ g IgG in 20 μ L volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at -20°C .

REACTIVITY: This antibody reacts with human α -tubulin on western blotting, Immunoprecipitation, and Immunocytochemistry. The reactivity to mouse, rat, hamster and chicken α -tubulin was confirmed by Western blotting.

APPLICATIONS:

Western blotting; 2 μ g/mL for chemiluminescence detection system

Immunoprecipitation; 5 μ g/200 μ L of cell extract from 2×10^6 cells

Immunohistochemistry; Not tested

Immunocytochemistry; 2 μ g/mL

Flow cytometry; Not tested

Detailed procedure is provided in the following **PROTOCOLS**.

SPECIES CROSS REACTIVITY:


Species	Human	Mouse	Rat	Hamster	Chicken
Cells	HeLa	NIH/3T3	NRK	CHO	MuH1
Reactivity on WB	+	+	+	+	+

REFERENCES:

- 1) Heald, R., and Nogales, E., *J. Cell Sci.* **115**, 3-4 (2002)
- 2) Hall, J. L., and Cowan, N. J., *Nucleic Acids Res.* **13**, 207-223 (1985)

INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.



Western blot analysis of α -Tubulin in HeLa (1), NIH/3T3 (2), PC12 (3), CHO (4) and MuH1 (5) using M175-3.
Sample volume: 2 μ g per lane

PROTOCOLS:


SDS-PAGE & Western Blotting

- 1) Wash cells (approximately 1×10^7 cells) 3 times with PBS and resuspend them in 10 volumes of cold Lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% NP-40) containing protease inhibitors at appropriate concentrations. Incubate it at 4°C with rotating for 30 minutes; thereafter, briefly sonicate the mixture (up to 10 seconds).
- 2) Centrifuge the tube at $12,000 \times g$ for 10 minutes at 4°C and transfer the supernatant to another tube. Measure the protein concentration of the supernatant and add the cold Lysis buffer to make 0.2 mg/mL solution.
- 3) Mix the sample with equal volume of Laemmli's sample buffer.
- 4) Boil the samples for 3 minutes and centrifuge. Load 20 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
- 5) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm^2 for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 6) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature, or overnight at 4°C .
- 7) Incubate the membrane for 1 hour at room temperature

with primary antibody diluted with PBS (pH 7.2) containing 1% skimmed milk as suggested in the **APPLICATIONS**. (The concentration of antibody will depend on the conditions.)

- 8) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 9) Incubate the membrane with 1:10,000 HRP-conjugated anti-mouse IgG (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 10) Wash the membrane with PBS-T (5 minutes x 3 times).
- 11) Wipe excess buffer off the membrane, and incubate membrane with appropriate chemiluminescence reagent for 1 minute.
- 12) Remove extra reagent from the membrane by dabbing with a paper towel, and seal it in plastic wrap.
- 13) Expose the membrane onto an X-ray film in a dark room for 5 minutes. Develop the film under usual settings. The conditions for exposure and development may vary.

(Positive controls for Western blotting; HeLa, NIH/3T3, PC12, CHO, MuH1)




Immunoprecipitation of α -Tubulin from HeLa with mouse IgG2a isotype control (1) or M175-3 (2). After immunoprecipitated with the antibody, immunocomplexes were resolved on SDS-PAGE and immunoblotted with anti- α -Tubulin polyclonal antibody (MBL; Code no. PM054).

Immunoprecipitation

- 1) Wash cells (approximately 1×10^7 cells) 3 times with PBS and resuspend them in 1 mL of cold Lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.05% NP-40) containing protease inhibitors at appropriate concentrations. Incubate it at 4°C with rotating for 30 minutes; thereafter, briefly sonicate the mixture (up to 10 seconds).
- 2) Centrifuge the tube at 12,000 x g for 10 minutes at 4°C and transfer the supernatant to another fresh tube.
- 3) Add primary antibody as suggested in the **APPLICATIONS** into 200 μ L of the supernatant. Mix well and incubate with gentle agitation for 60-120 minutes at 4°C. Add 20 μ L of 50% protein A agarose beads resuspended in the cold IP buffer (10 mM Tris-HCl pH 8.0, 500 mM NaCl, 0.1% NP-40). Mix well and incubate with gentle agitation for 60 minutes at 4°C.
- 4) Wash the beads 3-5 times with the cold Lysis buffer (centrifuge the tube at 2,500 x g for 10 seconds).

- 5) Resuspend the beads in 20 μ L of Laemmli's sample buffer, boil for 3-5 minutes, and centrifuge for 5 minutes. Use 20 μ L/lane for the SDS-PAGE analysis. (See **SDS-PAGE & Western blotting**.)

(Positive control for Immunoprecipitation; HeLa)



Immunocytochemical detection of α -Tubulin in HeLa using M175-3.

Green: anti- α -Tubulin
Blue: DAPI counterstain

Immunocytochemistry

- 1) Culture the cells in the appropriate condition on a glass slide. (for example, spread 1×10^4 cells for one slide, then incubate in a CO₂ incubator for one night.)
- 2) Wash the glass slide 2 times with PBS.
- 3) Fix the cells by immersing the slide in PBS containing 4% paraformaldehyde for 10 minutes at room temperature.
- 4) Wash the glass slide 3 times with PBS.
- 5) Immerse the slide in PBS containing 0.2% Triton X-100 for 10 minutes at room temperature.
- 6) Wash the glass slide 2 times with PBS.
- 7) Add the primary antibody diluted with PBS containing 2% FCS as suggested in the **APPLICATIONS** onto the cells and incubate for 30 minutes at room temperature (Optimization of antibody concentration or incubation condition are recommended if necessary.)
- 8) Wash the glass slide 2 times with PBS.
- 9) Add 100 μ L of 1:500 Alexa Fluor[®] 488 conjugated anti-mouse IgG (Invitrogen; code no. A11001) diluted with PBS onto the cells. Incubate for 30 minutes at room temperature. Keep out light by aluminum foil.
- 10) Wash the glass slide 2 times with PBS.
- 11) Counter stain with DAPI for 5 minutes at room temperature.
- 12) Wash the glass slide 2 times with PBS.
- 13) Wipe excess liquid off the slide but take care not to touch the cells. Never leave the cells to dry.
- 14) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; HeLa)

RELATED PRODUCTS:

- M175-3 Anti- α -Tubulin mAb (2F9)
M175-A48 Anti- α -Tubulin mAb-Alexa Fluor[®] 488 (2F9)
M175-A59 Anti- α -Tubulin mAb-Alexa Fluor[®] 594 (2F9)
M175-A64 Anti- α -Tubulin mAb-Alexa Fluor[®] 647 (2F9)
PM054 Anti- α -Tubulin pAb
PM054-7 Anti- α -Tubulin pAb-HRP-Direct
M176-3 Anti-EEA1 mAb (3C10)
M176-A48 Anti-EEA1 mAb-Alexa Fluor[®] 488 (3C10)
M176-A59 Anti-EEA1 mAb-Alexa Fluor[®] 594 (3C10)
M176-A64 Anti-EEA1 mAb-Alexa Fluor[®] 647 (3C10)
PM062 Anti-EEA1 pAb
M178-3 Anti-Calnexin mAb (4F10)
M178-A48 Anti-Calnexin mAb-Alexa Fluor[®] 488 (4F10)
M178-A59 Anti-Calnexin mAb-Alexa Fluor[®] 594 (4F10)
M178-A64 Anti-Calnexin mAb-Alexa Fluor[®] 647 (4F10)
PM060 Anti-Calnexin pAb
M181-3 Anti-KDEL mAb (1D5)
PM059 Anti-KDEL pAb
M179-3 Anti-GM130 mAb (5G8)
M179-A48 Anti-GM130 mAb-Alexa Fluor[®] 488 (5G8)
M179-A59 Anti-GM130 mAb-Alexa Fluor[®] 594 (5G8)
M179-A64 Anti-GM130 mAb-Alexa Fluor[®] 647 (5G8)
PM061 Anti-GM130 pAb
PM063 Anti-COX4 pAb
PM064 Anti-Lamin B1 pAb
- D115-3 Anti-CENP-A (Human) mAb (3-19)
PD030 Anti-CENP-C (Human) pAb
K0171-3 Anti-CENP-E (Human) mAb (1H12)
PD031 Anti-CENP-H (Human) pAb
PD032 Anti-CENP-I (hMis6) (Human) pAb
D282-3 Anti-CENP-K (ICEN37) (Human) mAb (46F1)
PD018 Anti-CENP-K (ICEN37) (Human) pAb
D283-3 Anti-CENP-L (ICEN33) (Human) mAb (27E10)
D284-3 Anti-CENP-M (ICEN39) (Human) mAb (23F6)
D285-3 Anti-CENP-N (ICEN32) (Human) mAb (22F4)
PD020 Anti-CENP-O (Chicken) pAb
D286-3 Anti-CENP-T (ICEN22) (Human) mAb (42F10)
PD019 Anti-CENP-50 (Human) pAb
D288-3 Anti-MgcRacGAP (Human) mAb (5G5)
- PM036 Anti-LC3 pAb [WB, IP, IC, IHC, FCM]
M152-3 Anti-LC3 mAb (4E12) [WB, IP, IC, FCM, EM]
M186-3 Anti-LC3 mAb (8E10) [WB]
PD014 Anti-LC3 pAb [WB]
PD015 Anti-LC3 pAb [IC]
PM046 Anti-LC3 pAb [WB, IC]
M115-3 Anti-LC3 mAb (51-11) [WB]
PM040 Anti-Atg16L pAb
M150-3 Anti-Atg16L mAb (1F12)

- WB: Western blotting
IP: Immunoprecipitation
IC: Immunocytochemistry
IHC: Immunohistochemistry
FCM: Flow cytometry
EM: Immuno-electron microscopy

Other related antibodies and kits are also available.
Please visit our website at <http://ruo.mbl.co.jp/>