

For Research Use Only.
Not for use in diagnostic procedures.



Anti-FcεR1γ (FcRγ) (Mouse) mAb -Alexa Fluor[®] 594

CODE No. M191-A59

CLONALITY Monoclonal
CLONE 1D6
ISOTYPE Mouse IgG1 κ
QUANTITY 100 μL, 1 mg/mL

SOURCE Purified IgG from hybridoma supernatant
FORMURATION PBS containing 1% BSA and 0.09% NaN₃

*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.

STORAGE This antibody solution is stable for one year from the date of purchase when stored at 4°C.

APPLICATIONS-CONFIRMED

Immunocytochemistry 1 μg/mL

SPECIES CROSS REACTIVITY

Species	Human	Mouse	Rat	Hamster
Cell	Not tested	RAW264	Not tested	Not Tested
Reactivity		+		

Entrez Gene ID 14127 (Mouse)

REFERENCES

- 1) Yamasaki, S., *et al.*, *Nat. Immunol.* **9**, 1179-1188 (2008)
- 2) Cao, L., *et al.*, *J.Immunol.* **179**, 5864-5876 (2007)
- 3) Sato, K., *et al.*, *J. Biol. Chem.* **281**, 38854-38866 (2006)
- 4) Ra, C., *et al.*, *J. Biol. Chem.* **264**, 15323-15327 (1989)

For more information, please visit our web site <https://ruo.mbl.co.jp/>

LABEL LICENSES:

This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co., LTD, and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by or licensed to LIFE TECHNOLOGIES Corporation. For information on purchasing a license to this product for any other use, contact LIFE TECHNOLOGIES Corporation, Molecular Probes Detection Technologies, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc.


RELATED PRODUCTS

- PM068 Anti-FcεR1γ (FcRγ) (Mouse) pAb
- M191-3 Anti-FcεR1γ (FcRγ) (Mouse) mAb (1D6)
- M191-A48 Anti-FcεR1γ (FcRγ) (Mouse) mAb
-Alexa Fluor® 488 (1D6)
- M191-A64 Anti-FcεR1γ (FcRγ) (Mouse) mAb
-Alexa Fluor® 647 (1D6)
- D292-3 Anti-Mincle (Mouse) mAb (4A9)
- D303-3 Anti-FcμR (Toso/Faim3) (Mouse) mAb (#4B5)

Immunocytochemistry

- 1) Spread the cells on a glass slide, then incubate in a CO₂ incubator for one night.
- 2) Remove the culture supernatant by careful aspiration.
- 3) Fix the cells by immersing the slide in 4% paraformaldehyde (PFA)/PBS for 10 min. at room temperature (20~25°C).
- 4) Prepare a wash container such as a 500 mL beaker with a magnetic stirrer. Then wash the fixed cells on the glass slide by soaking the slide with a plenty of PBS in the wash container for 5 min. Take care not to touch the cells. Repeat another wash once more.
- 5) Immerse the slide in 0.2% Triton X-100/PBS for 10 min. at room temperature.
- 6) Wash the slide 2 times in PBS for 5 min. each.
- 7) Add Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell and incubate for 5 min. at room temperature.
- 8) Add the primary antibody diluted with 2% fetal calf serum (FCS)/PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hr. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 9) Wash the slide 2 times in PBS for 5 min. each.
- 10) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 11) Counter stain with DAPI for 5 min. at room temperature.
- 12) Wash the slide 2 times in PBS for 5 min. each.
- 13) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; RAW264)



Immunocytochemical detection of

FcεR1γ (Mouse) in RAW264

Red: M191-A59

Green: DAPI