

**For Research Use Only.**  
**Not for use in diagnostic procedures.**



# Anti-CD36 (GPIV) (Human) mAb

**CODE No.** M232-3

**CLONALITY** Monoclonal  
**CLONE** 1-3  
**ISOTYPE** Mouse IgG2b  $\kappa$   
**QUANTITY** 100  $\mu$ g/100  $\mu$ L

**SOURCE** Purified IgG from hybridoma supernatant  
**IMMUNOGEN** Human CD36 expressed 293T transfectants  
**FORMULATION** PBS containing 50% Glycerol (pH 7.2). No preservative is contained.  
**STORAGE** This antibody solution is stable for one year from the date of purchase when stored at -20°C.

## APPLICATION-CONFIRMED

Flow cytometry 1  $\mu$ g/mL (final concentration)

## SPECIES CROSS REACTIVITY on FCM

Species	Human	Mouse	Rat	Hamster
Samples	HEL	Not tested	Not tested	Not tested
Reactivity	+			

**Entrez Gene ID** 948 (Human)

**REFERENCES**

- 1) Srikanthan, S., *et al.*, *J. Thromb. Haemost.* **12**, 1906-1917 (2014)
- 2) Welton, J. L., *et al.*, *Mol. Cell Proteomics* **9**, 1324-1338 (2010)
- 3) Admyre, C., *et al.*, *J. Immunol.* **179**, 1969-1978 (2007)

For more information, please visit our web site <http://ruo.mbl.co.jp/>



MEDICAL & BIOLOGICAL LABORATORIES CO., LTD.  
URL <http://ruo.mbl.co.jp/>  
e-mail [support@mbi.co.jp](mailto:support@mbi.co.jp), TEL 052-238-1904

**RELATED PRODUCTS**Antibodies

M232-3	Anti-CD36 (GPIV) (Human) mAb (1-3)
M232-6	Anti-CD36 (GPIV) (Human) mAb-Biotin (1-3)
D281-6	Anti-CD61 (GPIIIa) (Human) mAb-Biotin (T74)
D281-3	Anti-CD61 (GPIIIa) (Human) mAb (T74)
D281-A48	Anti-CD61 (GPIIIa) (Human) mAb -Alexa Fluor <sup>®</sup> 488 (T74)
D281-A64	Anti-CD61 (GPIIIa) (Human) mAb -Alexa Fluor <sup>®</sup> 647 (T74)
MEX001-3	Anti-CD9 mAb (A100-4)
MEX001-4	Anti-CD9 mAb-FITC (A100-4)
MEX001-6	Anti-CD9 mAb-Biotin (A100-4)
MEX001-12	Anti-CD9 mAb-ALP (A100-4)
MEX002-3	Anti-CD63 (LAMP-3) mAb (C047-1)
MEX002-4	Anti-CD63 (LAMP-3) mAb-FITC (C047-1)
MEX002-6	Anti-CD63 (LAMP-3) mAb-Biotin (C047-1)
MEX002-12	Anti-CD63 mAb-ALP (C047-1)
MEX003-3	Anti-CD81 (TAPA1) mAb (A103-10)
MEX003-4	Anti-CD81 (TAPA1) mAb-FITC (A103-10)
MEX003-6	Anti-CD81 (TAPA1) mAb-Biotin (A103-10)
MEX003-12	Anti-CD81 (TAPA1) mAb-ALP (A103-10)
MEX004-6	Anti-CD326 (EpCAM) mAb-Biotin (B8-4)
D252-3	Anti-CD9 (Human) mAb (10H6)
D252-5	Anti-CD9 (Human) mAb-PE (10H6)
D131-3	Anti-CD9 (Mouse) mAb (JF9)
D131-4	Anti-CD9 (Mouse) mAb-FITC (JF9)
D263-3	Anti-CD63 (LAMP-3) (Mouse) mAb (R5G2.1)
D082-3	Anti-CD151 (SFA-1) (Human) mAb (SFA1.2B4)
D082-5	Anti-CD151 (SFA-1) (Human) mAb-PE (SFA1.2B4)
D050-3	Anti-CD29 (Integrin $\beta$ 1) (Human) mAb (AG89)
D050-5	Anti-CD29 (Integrin $\beta$ 1) (Human) mAb-PE (AG89)
D276-3	Anti-CD36 (GPIV) (Human) mAb (GS95)
D276-A48	Anti-CD36 (GPIV) (Human) mAb -Alexa Fluor <sup>®</sup> 488 (GS95)
D276-A64	Anti-CD36 (GPIV) (Human) mAb -Alexa Fluor <sup>®</sup> 647 (GS95)
D269-3	Anti-EpCAM (CD326) (Mouse) mAb (2-17-F1)
K0142-3	Anti-PSMA (Human) mAb (107-1A4)
K0142-4	Anti-PSMA (Human) mAb-FITC (107-1A4)
K0142-5	Anti-PSMA (Human) mAb-PE (107-1A4)
K0142-6	Anti-PSMA (Human) mAb-Biotin (107-1A4)
M077-3	Mouse IgG2b (isotype control) (3D12)

Kits

MEX-SA	ExoCap <sup>™</sup> Streptavidin Kit
MEX-SA123	ExoCap <sup>™</sup> Streptavidin CD9/CD63/CD81 Set-Biotin
MEX-E	ExoCap <sup>™</sup> Nucleic Acid Elution Buffer
MEX1001	ExoDiluent for Immunoassay

Equipment

3190	Magnetic Rack
------	---------------


Other related antibodies and kits are also available.

Please visit our website at <http://ruo.mbl.co.jp/>

**Flow cytometric analysis**

- 1) Wash the cells ( $3 \times 10^5$  cells/sample) 3 times with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 2) Add 10  $\mu$ L of Clear Back (MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 10 min. at room temperature.
- 3) Add 30  $\mu$ L of primary antibody diluted with washing buffer as suggested in the **APPLICATION**. Mix well and incubate for 30 min. at room temperature.
- 4) Wash the cells 1 time with 1 mL of washing buffer.
- 5) Add 30  $\mu$ L of 1:500 Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488 (Thermo Fisher Scientific; A-11001) diluted with washing buffer. Mix well and incubate for 15 min. at room temperature.
- 6) Wash the cells 1 time with 1 mL of the washing buffer.
- 7) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; HEL)



***Flow cytometric detection of human CD36 (GPIV) expressed on HEL cells***

Open: Anti-CD36 (GPIV) (Human) mAb (M232-3)

Closed: Mouse IgG2b (isotype control) (M077-3)