

Normal Guinea Pig IgG

CODE No.	PM067
CLONALITY	Polyclonal
QUANTITY	100 µL, 1 mg/mL
SOURCE	Purified IgG from normal guinea pig serum using protein A agarose.
REACTIVITY	No specific reaction was detected on immunoprecipitation and flow cytometry.
FORMULATION	1 mg/mL in PBS containing 50% glycerol. No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATIONS-CONFIRMED

Immunoprecipitation

Flow cytometry

This antibody can be used as a negative isotypic control.
The concentration will depend on the conditions.

APPLICATION-REPORTED

Immunohistochemistry Reference 1) and 2)

REFERENCES	1) Murai, N., <i>et al.</i> , <i>PLoS One</i> 12 , e0186637 (2017) [IHC]
	2) Yamane, T., <i>et al.</i> , <i>PLoS One</i> 12 , e0176809 (2017) [WB, IHC]

For more information, please visit our web site <http://ruo.mbl.co.jp/>

RELATED PRODUCTS

Functional grade antibodies

M075-3M2 Mouse IgG1 (isotype control) (2E12)
M076-3M2 Mouse IgG2a (isotype control) (6H3)
M077-3M2 Mouse IgG2b (isotype control) (3D12)
M078-3M2 Mouse IgG3 (isotype control) (6A3)
M080-3M2 Rat IgG1 (isotype control) (1H5)
M081-3M2 Rat IgG2a (isotype control) (2H3)
M090-3M2 Rat IgG2b (isotype control) (3G8)

Purified antibodies


M075-3 Mouse IgG1 (isotype control) (2E12)
M075-4 Mouse IgG1 (isotype control)-FITC (2E12)
M075-5 Mouse IgG1 (isotype control)-PE (2E12)
M075-6 Mouse IgG1 (isotype control)-Biotin (2E12)
M075-11 Mouse IgG1 (isotype control)-Magnetic Beads (2E12)
M075-12 Mouse IgG1 (isotype control)-ALP (2E12)
M075-A48 Mouse IgG1 (isotype control)-Alexa Fluor[®] 488 (2E12)
M075-A64 Mouse IgG1 (isotype control)-Alexa Fluor[®] 647 (2E12)
M075-8 Mouse IgG1 (isotype control)-Agarose (2E12)
M076-3 Mouse IgG2a (isotype control) (6H3)
M076-4 Mouse IgG2a (isotype control)-FITC (6H3)
M076-5 Mouse IgG2a (isotype control)-PE (6H3)
M076-6 Mouse IgG2a (isotype control)-Biotin (6H3)
M076-11 Mouse IgG2a (isotype control)-Magnetic Beads (6H3)
M076-12 Mouse IgG2a (isotype control)-ALP (6H3)
M076-A48 Mouse IgG2a (isotype control)-Alexa Fluor[®] 488 (6H3)
M076-A64 Mouse IgG2a (isotype control)-Alexa Fluor[®] 647 (6H3)
M077-3 Mouse IgG2b (isotype control) (3D12)
M077-4 Mouse IgG2b (isotype control)-FITC (3D12)
M077-5 Mouse IgG2b (isotype control)-PE (3D12)
M077-6 Mouse IgG2b (isotype control)-Biotin (3D12)
M077-11 Mouse IgG2b (isotype control)-Magnetic Beads (3D12)
M077-12 Mouse IgG2b (isotype control)-ALP (3D12)
M077-A48 Mouse IgG2b (isotype control)-Alexa Fluor[®] 488 (3D12)
M077-A64 Mouse IgG2b (isotype control)-Alexa Fluor[®] 647 (3D12)
M078-3 Mouse IgG3 (isotype control) (6A3)
M078-4 Mouse IgG3 (isotype control)-FITC (6A3)
M078-6 Mouse IgG3 (isotype control)-Biotin (6A3)
M079-3 Mouse IgM (isotype control) (7E10)
M080-3 Rat IgG1 (isotype control) (1H5)
M080-4 Rat IgG1 (isotype control)-FITC (1H5)
M080-5 Rat IgG1 (isotype control)-PE (1H5)
M080-A48 Rat IgG1 (isotype control)-Alexa Fluor[®] 488 (1H5)
M080-A64 Rat IgG1 (isotype control)-Alexa Fluor[®] 647 (1H5)
M081-3 Rat IgG2a (isotype control) (2H3)
M081-4 Rat IgG2a (isotype control)-FITC (2H3)
M081-5 Rat IgG2a (isotype control)-PE (2H3)
M081-A48 Rat IgG2a (isotype control)-Alexa Fluor[®] 488 (2H3)
M081-A64 Rat IgG2a (isotype control)-Alexa Fluor[®] 647 (2H3)
M081-8 Rat IgG2a (isotype control)-Agarose (2H3)
M081-11 Rat IgG2a (isotype control)-Magnetic Beads (2H3)
M082-3 Rat IgG2c (isotype control) (6E12)
M082-4 Rat IgG2c (isotype control)-FITC (6E12)
M090-3 Rat IgG2b (isotype control) (3G8)
M090-4 Rat IgG2b (isotype control)-FITC (3G8)
M090-5 Rat IgG2b (isotype control)-PE (3G8)
M090-A48 Rat IgG2b (isotype control)-Alexa Fluor[®] 488 (3G8)
M090-A64 Rat IgG2b (isotype control)-Alexa Fluor[®] 647 (3G8)
PM035 Normal Rabbit IgG (polyclonal)

PM035-8 Normal Rabbit IgG-Agarose (polyclonal)
M189-3 Syrian Hamster IgG (isotype control)
M199-3 Armenian Hamster IgG (isotype control)
PM084 Normal Chicken IgY (polyclonal)
PM084-4 Normal Chicken IgY-FITC (polyclonal)
PM089 Normal Sheep IgG (polyclonal)
PM094 Normal Goat IgG (polyclonal)

Other related antibodies and kits are also available.
Please visit our website at <http://ruo.mbl.co.jp/>

Immunoprecipitation

- 1) Wash 1×10^7 cells 2 times with PBS and resuspend them with 1 mL of ice-cold Lysis buffer [50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.05% NP-40] containing appropriate protease inhibitors. Incubate it on ice for 15 min., thereafter, sonicate briefly (up to 15 sec.).
- 2) Centrifuge the tube at 12,000 x g for 5 min. at 4°C and transfer the supernatant to another tube.
- 3) Add the isotype control antibody at the equal amount of the antibody for immunoprecipitation to the supernatant. Vortex briefly and incubate with gentle agitation for 60-120 min. at 4°C.
- 4) Mix 20 μ L of 50% protein A agarose beads slurry resuspended in 400 μ L of IP buffer [10 mM Tris-HCl (pH 8.0), 500 mM NaCl, 0.1% NP-40] with primary antibody. Incubate with gentle agitation for 1 hr. at room temperature.
- 5) Wash the beads 3 times with 1 mL of IP buffer.
- 6) Add 300 μ L of cell lysate (prepared sample of step 2), then incubate with gentle agitation for 1 hr. at room temperature.
- 7) Centrifuge the tube at 2,500 x g for 10 sec. Carefully remove and discard the supernatant.
- 8) Resuspend the beads with 1 mL of Lysis buffer.
- 9) Centrifuge the tube at 2,500 x g for 10 sec. Carefully remove and discard the supernatant.
- 10) Repeat Steps 8)-9) 5 times.
- 11) Resuspend the beads in 20 μ L of Laemmli's sample buffer, boil for 3 min. and centrifuge for 5 min.
- 12) Load 10 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel for electrophoresis.
- 13) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 14) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 15) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 16) Wash the membrane with PBS (5 min. x 3 times).
- 17) Incubate the membrane with Anti-IgG (Rabbit) pAb-HRP (MBL; code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 18) Wash the membrane with PBS (5 min. x 3 times).
- 19) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min.
- 20) Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 21) Expose to an X-ray film in a dark room for 30 sec. Develop the film as usual. The condition for exposure and development may vary.



Immunoprecipitation of mouse p62 from NIH/3T3 cells


Lane 1: IP with Normal Guinea Pig IgG (PM067)

Lane 2: IP with Anti-p62 C-terminal pAb (MBL; code no. PM066)

Immunoblotted with Anti-p62 (SQSTM1) pAb (MBL; code no. PM045)

Flow cytometric analysis for adherent cells

- 1) Detach the cells from culture dish.
- 2) Wash the cells 3 times with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 3) Add 200 μ L of 4% paraformaldehyde (PFA) to the cell pellet after tapping. Mix well, then fix the cells for 10 min. at room temperature.
- 4) Wash the cells 2 times with 1 mL of washing buffer.
- 5) Add 200 μ L of PBS containing 100 μ g/mL Digitonin to the cell pellet after tapping. Mix well, then permeabilize the cells for 10 min. at room temperature.
- 6) Wash the cells 2 times with 1 mL of washing buffer.
- 7) Resuspend the cells with washing buffer (5×10^6 cells/mL).
- 8) Add 100 μ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 min. at room temperature (20~25°C). Remove the supernatant by careful aspiration.
- 9) Add 20 μ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 min. at room temperature.
- 10) Add the isotype control antibody at the concentrations comparable to those of the specific antibody of interest. Mix well and incubate for 30 min. at room temperature.
- 11) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration.
- 12) Add FITC-conjugated anti-Guinea Pig IgG antibody diluted with the washing buffer. Mix well and incubate for 30 min. at room temperature.
- 13) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration.
- 14) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.



Flow cytometric analysis of p62 in HeLa

Closed: Secondary antibody alone

Open: Normal Guinea Pig IgG (PM067) or Anti-p62 C-terminal pAb (MBL; code no. PM066)