

**MONOCLONAL ANTIBODY**

# Anti-RECK (Human) mAb

Code No.	Clone	Subclass	Quantity	Concentration
W050-3	2C3D1F	Mouse IgG2a $\kappa$	100 $\mu$ L	1 mg/mL

**BACKGROUND:** Reversion-inducing cysteine-rich protein with Kazal motifs (RECK) functions as a tumor and metastasis suppressor through inhibiting proteolytic activity of matrix metalloproteinase-9. While RECK is expressed in a wide variety of tissues and normal cell lines, its expression is weak or absent in various cancer types including colon cancer, breast cancer and glioma. Over expression of RECK in tumor cell lines doesn't affect their growth but significantly suppresses matrix invasion and metastatic activity.

**SOURCE:** This antibody was purified from hybridoma culture supernatant by Protein A affinity column chromatography.

**IMMUNOGEN:** Human RECK expressed Ba/F3 transfectants generated from SST-REX (signal sequence trap by retrovirus-mediated expression screening).

**FORMULATION:** 100  $\mu$ g IgG in 100  $\mu$ L volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at  $-20^{\circ}\text{C}$ .

**REACTIVITY:** This antibody reacts with human RECK on Flow cytometry.

**APPLICATIONS:**

- Flow cytometry; 1-10  $\mu$ g/mL
- Western blotting; Not tested
- Immunoprecipitation; Not tested
- Immunohistochemistry; Not tested
- Immunocytochemistry; Not tested

Detailed procedure is provided in the following **PROTOCOL**.

**INTENDED USE:**

For Research Use Only. Not for use in diagnostic procedures.

**Entrez Gene ID:**


8434 (Human)

**REFERENCES:**

- 1) Oh, J., *et al.*, *Cell* **107**, 789-800 (2001)
- 2) Kojima, T and Kitamura, T. *Nat. Biotechnol.* **17**, 487-490 (1999)
- 3) Takahashi, C., *et al.*, *PNAS* **95**, 13221-13226 (1998)

**SPECIES CROSS REACTIVITY:**

Species	Human	Mouse	Rat	Hamster
Cells	Transfectant	Not tested	Not tested	Not tested
Reactivity on FCM	+			



**Flow cytometric analysis of human RECK expression on Ba/F3 transfectant.** Open histograms indicate the reaction of isotypic control to the cells. Shaded histograms indicate the reaction of W050-3 to the cells.

**PROTOCOL:**

**Flow cytometric analysis for floating cells**

We usually use Fisher tubes or equivalents as reaction tubes for all steps described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.05%  $\text{NaN}_3$ ].
- 2) Resuspend the cells with washing buffer ( $2.5 \times 10^6$  cells/mL).
- 3) Add 200  $\mu$ L of cell suspension into each tube. And centrifuge at 500 x g for 1 minute at room temperature ( $20\sim 25^{\circ}\text{C}$ ). Remove supernatant by careful decantation.
- 4) Add 20  $\mu$ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 minutes at room

temperature.

- 5) Add 50  $\mu$ L of the primary antibody at the concentration as suggest in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful decantation.
- 7) Add 50  $\mu$ L of 1:200 anti-mouse IgG-PE (Beckman Coulter; code no. IM0855) diluted with the washing buffer. Mix well and incubate for 30 minutes at room temperature.
- 8) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful decantation.
- 9) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; transfectant)

#### **RELATED PRODUCTS:**

D049-3	Anti-RECK (Human) mAb
W005-3	Anti-BTN2A1 (Human) mAb
W008-3	Anti-Carboxypeptidase D (Human) mAb
W011-3	Anti-Dystroglycan (Human) mAb
W017-3	Anti-EphA2 (Human) mAb
W029-3	Anti-IGFBP1 (Human) mAb
W050-3	Anti-RECK (Human) mAb
W086-3	Anti-LYPD3 (C4.4A) (Human) mAb
W089-3	Anti-C11orf24 (Human) mAb
W124-3	Anti-GPR56 (Human) mAb
W125-3	Anti-GPR56 (Human) mAb
W194-3	Anti-FAM171A1 (Human) mAb
W253-3	Anti-Glypican 1 (Human) mAb
W357-3	Anti-CD105 (Endoglin) (Human) mAb
W358-3	Anti-CD300A (Human) mAb
W359-3	Anti-CD300C (Human) mAb
M076-3	Mouse IgG2a (isotype control)
MTG-001	Clear Back (Human Fc receptor blocking reagent)